

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DISCOVERY OF COMET e 1898 (PERRINE).

This comet was discovered in the morning of June 15th, in the constellation *Camelopardalis*. At 23^h 22^m 34^s G. M. T. of June 14th, the comet's position was $\alpha = 3^h$ 29^m os 99, $\delta = +58^\circ$ 35′ 22.″3. It was then moving slowly south and more rapidly east. The elements as computed from the first three observations will be found elsewhere in this number of the *Publications*. These elements bear considerable resemblance to those of the comet 1785 I, especially with regard to ω and ι , the resemblance being closer than was the case with the comet b 1898, pointed out in No. 62 of these *Publications*. Comet e 1898 appears to be another member of this same family.

At the time of discovery the comet was not so bright as Comet δ , its brightness being estimated at about 10 or a little fainter. It has been steadily increasing in brightness and is now estimated to be 9 or $9\frac{1}{2}$ magnitude, brighter than Comet δ . A rather sharp nucleus has developed within the past week and is now estimated to be about 13th magnitude.

Comet e has been moving in the same general direction as Comet b, and as its geocentric motion has been much more rapid it has overtaken the latter and passed it in both co-ordinates. On the morning of June 27th, the two comets were within about one-quarter degree of each other, both visible in the lowest power field of the 12-inch telescope.

Comet *e* is moving rapidly south and east, and hence its location is becoming more unfavorable for observation. Towards the middle of August it will pass the Sun going south, after which it will soon be lost to northern observers. Owing to the resemblance already pointed out, it seems important to observe it as long as possible.

C. D. Perrine.

Mt. Hamilton, Cal., 1898, June 29.

ERRATA IN STAR CATALOGUES.

Lalande 31379. The declination of this star appears to be in error by 1'. A micrometer comparison with Radcliffe_{3'} 4514 on June 2d indicates that the N. P. D. of Lalande 31379 should be 100° 25′ 39″.8.

WEISSE'S Bessel XVIII, 1327 appears to be in error in declination by 2'. A micrometer comparison with 1323 on June 2d shows that the declination of 1327 should be -13° 44' 10".5.

C. D. PERRINE.

Mt. Hamilton, Cal., June 7, 1898.